D-1

TECTONICS OF THE EASTERN MEDITERRANEAN

XAVIER LEPICHON, CLAUDE RANGIN and NICOLAS CHAMOT-ROOKE
 CNRS, Laboratoire de Géologie, Ecole Normale Supérieure
 24 rue Lhomond, 75231 Paris, France

There has been a considerable amount of studies of the lands bordering the Eastern Mediterranean basin but the basin itself is still relatively poorly known. The tectonic analysis of this basin is difficult to do because a great part of it has disappeared within a double system of subduction zones, the Hellenic one that is still active and the Calabrian one that stopped being active less than one million years ago. To discuss the tectonic evolution of the Eastern Mediterranean basin, it is first necessary to evaluate the effect of the present kinematics and then to reconstruct the geometry of the basin prior to subduction. We now are able to precisely evaluate using geodetic methods these present kinematics that are dominated by the lateral extrusion of Anatolia and the N/S extension of Aegea. We have the possibility to reconstruct rather confidently the middle Miocene shape of the basin that was then about twice larger than today. It is much more difficult to reconstruct the geometry of the basin prior to Miocene. Important guides for such reconstructions are the present structures of the basins, of the accretionary wedges that cover half of them and of their margins. Based in great part on work that has been obtained by our own laboratory, we summarize our present state of knowledge about these three topics:

1) *The crustal structure of the basins.* There are significant differences between the western (Ionian) and eastern (Levantine) basins. The structure of the Ionian basin is better known. It is clearly oceanic. The sedimentary cover is well stratified and the main layers may be followed over the whole basin. The Ionian basin was affected by active shortening prior to Messinian, and most probably during late middle Miocene. The structure of the Levantine basin is more difficult to characterize due to the much thicker sedimentary cover. It is compatible either with an oceanic crust or with a thinned continental crust.

2) *The structure and development of the Mediterranean and Calabrian wedges.* The Mediterranean ridge is the accretionary complex formed since Middle Miocene time by the Hellenic subduction. We have proposed that the backstop against which this accretionary complex is formed consists of Hellenic nappes that came from the Aegea during lower Miocene. The structure of the Calabrian wedge is less well known but appears to be mostly related to the post-middle Miocene Calabrian subduction that now appears to be inactive. We have identified a major still active transform fault that limits the Calabrian wedge on its western side and that appears to be linked to the formation of the Etna. We relate its activity to the progressive verticalization of the Calabrian (Tyrrhenian) slab.

2) *The ages and developments of the southern (African) African and northern (European) margins.* We argue that the age of the African margin of the Levantine basin is Triassic whereas it is middle Cretaceous for the Ionian basin. A phase of subsidence of this age is well known over several plateaus, seamounts and ridges there. Middle Cretaceous is probably the age of the formation of the oceanic crust of the Ionian basin. We examine the northern European margin north of the Hellenic trench and discuss possible evolutionary schemes for them.